



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Cerastes cerastes* (Linnæus). Two adults.  
*Ptyodactylus hasselquistii* (Donndorff). One adult.  
*Tarentola annularis* (G. St. Hilaire). Four adults.  
*Agama flavimaculata* (Rüppell). Adult.  
*Agama pallida* Reuss. Three adults.  
*Uromastix ornatus* Rüppell. One adult. This species  
 apparently not previously recorded definitely  
 from Egypt. Back uniform in color.  
*Varanus griseus* (Daudin). Adult.  
*Varanus niloticus* (Linnæus). Adult.  
*Acanthodactylus boskianus* (Daudin). One.  
*Scincus scincus* (Linnæus). Six, mostly adults.

HENRY W. FOWLER,  
*Philadelphia, Pa.*

## AMBLYSTOMA OF LONG ISLAND

Three species of *Amblystoma* (*A. punctatum*,  
*A. opacum*, *A. tigrinum*) have been listed as being  
 found in the vicinity of New York City.\* Of these  
 only *A. tigrinum* is mentioned, without specific  
 data, as occurring on Long Island. Observations  
 by the writer and material in the Museum of the  
 Brooklyn Institute of Arts and Sciences show the  
 following Long Island records:

### *AMBLYSTOMA PUNCTATUM* (Linnæus).

(Spotted Salamander).

2 specimens Cold Spring Harbor, L. I. Aug., 1900.  
 —G. K. Cherrie and L. Ericson.

3 specimens Forest Park, Brooklyn. May, 1906.  
 Under stones and logs, damp soil.  
 —Geo. P. Engelhardt.

1 specimen Huntington, L. I. Aug., 1913.  
 In potato patch. —Miss Graeve.

---

\*Sherwood, William L. Salamanders found in the vicinity of New York City.  
 Linn. Soc. 1894-95. No. 7.

Ditmars, Raymond L. The Batrachians of the vicinity of New York City.  
 Amer. Mus. Journal, Vol. V, No. 4. Oct., 1905.

This species appears to be common on Long Island. Specimens are frequently brought to the Museum. Its favorite haunts are damp woods, often at some distance from streams and ponds.

*AMBLYSTOMA OPACUM* (Gravenhorst).

(Marbled Salamander).

1 specimen Mt. Sinai, L. I. Sept. 18, 1909.

—R. C. Murphy.

1 specimen Baldwin, L. I. July 25, 1911.

—Dwight Franklin.

My only experience in finding this species has been in the pine barrens near Lakehurst, N. J., at a place known as "Bone Hill," where, in company with Mr. William T. Davis, three specimens were captured, Aug., 1906, in a barrel sunk at one end of a small pool. This pool, though much reduced in summer, never runs dry, and the barrel formed an excellent trap which the salamanders could readily enter in spring, when the water is high, but from which they could not escape in the summer. For several succeeding seasons *A. opacum* was found in the barrel, but no other species. When visited during June, 1912, the barrel had collapsed, but the pool swarmed with larvæ 1 to 1½ inches long. Except during the breeding season, this species prefers dry or shady places, as already reported by Cope and others.

*AMBYSTOMA JEFFERSONIANUM* (Green).

1 specimen Montauk, L. I. May 31, 1913.

—Geo. P. Engelhardt.

The capture of this specimen is of unusual interest in furnishing the first record of the species from Long Island and, it appears, also from the vicinity of New York City. In general it can be

said that this salamander is northern in its distribution. Cope records it from West Virginia, Pennsylvania, Ohio, Wisconsin, New York, Vermont and Canada; De Kay from Albany and Peru, Clinton County, New York. The Long Island specimen was discovered under a board in a slightly moist depression near the road about half way between Montauk and Montauk Point, a locality abounding in swamps, ponds and small streams. It measures four inches in length, therefore not fully grown, and in color during life was above glossy black with numerous light-blue blotches and spots; below grayish to lead-colored. During four months of captivity the specimen fed sparingly on meal worms. In alcohol the ground color has changed to olive-brown and the blotches and spots have faded into buff. A wax cast, with natural colors restored, has been prepared.

*AMBLYSTOMA TIGRINUM* (Green).

(Tiger Salamander.)

Reported by William L. Sherwood from Rancocas, L. I., but this no doubt should read Rancocas, N. J., as there is no such place on Long Island.

Raymond L. Ditmars refers to this species as "occasionally found on Long Island."

No specimens or specific data are at hand. Information regarding this species, as well as other salamanders from Long Island, would be much appreciated.

GEORGE P. ENGELHARDT,  
*Brooklyn, N. Y.*